

2019 FORAGE MAIZE DESCRIPTIVE LIST

First choice varieties for favourable sites (varieties ranked in dry matter yield order)

VARIETY	Dry matter content at harvest (%)	Dry matter yield (t/ha)	Dry matter yield (% of 4 and 5 year varieties)	ME of fresh plant at harvest (MJ/kg dry matter)	ME yield of fresh plant at harvest (1000's MJ/ha)	Starch content of whole plant at harvest (%)	Starch yield of whole plant at harvest (t/ha)	Cell wall digestibility (%)	Early vigour (1-9)	Standing power at harvest (root lodging) (1-9)	Lodging (%) #	Brackling (%) #	Leaf senescence (1-9)	Eyespot rating (1-9) **	Year first listed
LG31218	30.5	18.7	105	11.60	217	29.5	5.51	59.6	7.6	8.0	1.9	0.0	7.5	5.4	2018
LG31211	30.3	18.5	105	11.60	215	28.4	5.26	60.1	7.6	8.0	2.0	0.0	7.5	4.6	2017
AVITUS KWS	35.5	18.5	104	11.63	215	36.1	6.66	57.3	7.6	8.2	0.9	0.7	5.7	5.7	2018
SPYCI CS	29.0	18.4	104	11.42	210	28.9	5.33	58.9	7.1	8.3	0.4	0.0	7.3	4.0	2016
GATSBY	31.8	18.4	104	11.51	212	31.4	5.78	58.2	7.2	8.3	0.2	0.7	7.4	5.4	2017
AUTENS KWS	33.5	18.4	104	11.48	211	32.9	6.06	57.4	7.5	7.9	2.3	0.5	6.1	5.0	2017
KOMPETENS	30.1	18.4	104	11.41	210	30.2	5.55	57.5	6.6	8.3	0.3	2.0	8.1	7.1	2016
AURELIUS KWS	33.0	18.4	104	11.44	210	32.4	5.95	57.1	7.6	8.2	1.0	0.8	6.2	4.6	2016
TIBERIO	29.6	18.4	104	11.20	206	28.6	5.26	56.8	7.4	8.1	1.4	1.5	7.6	6.3	2017
LG30209	30.9	18.3	103	11.54	212	30.3	5.55	58.8	7.5	8.1	1.3	0.2	7.3	3.3	2016
CRANBERRI CS	29.5	18.3	103	11.32	207	28.2	5.16	58.5	6.9	7.9	2.5	0.6	7.5	6.6	2018
BONNIE	32.3	18.2	103	11.66	213	32.1	5.86	59.4	7.2	8.3	0.5	0.4	7.5	5.8	2017
ABSALON	31.5	18.2	103	11.53	210	29.9	5.45	59.2	7.1	8.3	0.3	0.1	7.2	5.9	2017
EXPEDIA	30.7	18.2	103	11.60	211	31.3	5.69	59.0	7.1	8.3	0.3	0.2	7.3	5.7	2018
PINNACLE	34.3	18.1	102	11.75	213	35.9	6.50	58.2	7.3	8.2	0.9	0.7	7.2	6.6	2018
RODRIGUEZ KWS	32.9	18.1	102	11.60	209	34.8	6.29	57.9	6.9	8.3	0.1	0.5	6.8	3.8	2015
BELAMI CS	31.2	18.0	102	11.59	209	30.3	5.46	59.6	7.1	7.6	3.9	0.0	6.6	4.5	2017
ATRIUM	30.4	18.0	102	11.70	210	30.4	5.47	60.3	7.4	8.2	0.6	0.0	7.6	2.8	2010
ASGAARD	32.0	18.0	101	11.51	207	31.2	5.62	58.5	7.7	8.3	0.3	0.1	7.1	3.8	2014
AMBITION	34.2	17.9	101	11.61	208	34.6	6.20	57.8	7.4	8.3	0.4	0.0	6.8	6.3	2012
YUKON	33.9	17.9	101	11.48	206	35.4	6.34	56.4	7.1	8.3	0.4	0.2	6.6	5.0	2012
FIELDSTAR	33.0	17.9	101	11.54	206	33.1	5.92	57.9	7.3	8.3	0.1	0.4	7.1	6.5	2013
SY NORDICSTAR	32.3	17.9	101	11.50	205	32.3	5.76	58.5	7.4	8.2	0.9	0.3	7.3	7.5	2016
EMBLEM	32.0	17.8	101	11.54	206	33.2	5.91	58.3	7.3	8.3	0.3	0.5	6.7	5.7	2014
MADONIAS	33.3	17.7	100	11.51	204	34.1	6.04	57.1	7.1	8.2	0.6	1.1	6.2	5.5	2018
REASON	35.4	17.6	100	11.61	205	33.9	5.97	57.9	7.2	8.2	0.9	0.8	6.2	3.4	2016
SEVERUS	33.8	17.6	99	11.51	203	35.1	6.18	56.9	7.4	8.3	0.5	0.8	6.2	5.3	2013
GLORY	34.9	17.6	99	11.64	205	35.6	6.27	57.7	7.4	8.3	0.4	0.1	6.6	6.3	2013
SUNLITE	34.0	17.1	97	11.73	201	34.9	5.97	59.2	7.2	8.3	0.2	0.0	6.5	3.0	2014
SERGIO KWS	35.4	16.7	94	11.76	196	37.2	6.21	58.4	7.4	7.9	2.1	0.7	5.5	5.7	2014
RUBIERA KWS	36.5	16.6	94	11.80	196	37.3	6.20	58.3	7.4	7.9	2.2	0.6	5.1	5.4	2015
ACTIVATE	35.6	16.5	93	11.80	195	36.1	5.96	58.9	6.8	8.3	0.4	0.0	6.5	4.6	2012
EMMERSON	37.6	16.4	93	11.62	191	36.4	5.98	57.2	7.0	8.3	0.1	1.3	5.4	2.6	2015
AUGUSTUS KWS	36.8	16.4	92	11.75	192	36.9	6.03	58.1	7.3	8.0	1.8	1.6	5.0	6.0	2015
CITO KWS	37.4	15.9	90	11.82	188	37.6	6.00	58.5	7.2	8.3	0.4	0.4	4.7	5.4	2018
Trial statistics															
Mean of 4 & 5 year varieties	32.7	17.7	100	11.48	204	32.3	5.72	57.8	7.2	8.2	0.6	0.4	6.8	4.9	-
LSD (V.Cont)	0.82	0.46	2.6	0.123	-	1.34	-	0.64	0.29	-	-	-	0.45	-	-
SE (variety mean)	0.412	0.234	1.32	0.0618	-	0.678	-	0.325	0.145	-	-	-	0.226	-	-

On the 1-9 scales, high figures indicate that a variety shows the character to a high degree (e.g. high resistance)

Notes:

ME is calculated as D-value x 0.16

= values are re-transformed from the mean of the log data

Limited data for brackling % with some varieties only having 1 trials worth of data

** = Eyespot rating derived from inoculated nurseries rather than field scores

Data from the Forage Maize and Biogas Maize Descriptive Lists. Full data at www.bspb.co.uk



With thanks to the Maize Growers Association for part-funding the eyespot assessments

2019 FORAGE MAIZE DESCRIPTIVE LIST

Second choice varieties for favourable sites (varieties ranked in dry matter yield order)

VARIETY	Dry matter content at harvest (%)	Dry matter yield (t/ha)	Dry matter yield (% of 4 and 5 year varieties)	ME of fresh plant at harvest (MJ/kg dry matter)	ME yield of fresh plant at harvest (1000's MJ/ha)	Starch content of whole plant at harvest (%)	Starch yield of whole plant at harvest (t/ha)	Cell wall digestibility (%)	Early vigour (1-9)	Standing power at harvest (root lodging) (1-9)	Lodging (%) #	Brackling (%) #	Leaf senescence (1-9)	Eyespot rating (1-9) **	Year first listed
MILKSTAR	30.3	19.0	107	11.00	209	23.0	4.36	56.9	7.4	8.0	1.7	0.2	7.5	5.4	2018
CATHY	30.4	18.5	105	11.34	210	27.3	5.06	58.4	7.2	8.3	0.3	0.4	7.8	4.8	2015
AMBROSINI	30.0	18.5	104	11.14	206	28.8	5.33	56.0	7.3	8.3	0.3	0.0	7.5	5.2	2010
MAS 13M	30.1	18.3	103	10.86	199	27.2	4.98	54.8	6.8	8.2	0.6	0.4	7.3	5.2	2017
ES MARCO	29.3	18.1	102	11.18	202	29.6	5.36	56.0	6.4	8.2	0.9	0.7	7.9	6.9	2015
MONTY	32.2	18.1	102	11.47	207	30.2	5.46	58.5	6.9	8.3	0.4	0.4	7.5	4.5	2014
CODITANK	29.7	18.0	101	11.30	203	29.0	5.20	57.9	6.5	8.3	0.3	0.3	7.2	6.2	2015
PEREZ KWS	34.9	17.7	100	11.42	202	34.3	6.08	56.6	7.7	8.0	2.0	0.5	5.9	4.8	2015
RGT OXXGOOD	33.3	17.6	99	11.53	203	32.9	5.79	58.1	7.2	8.2	0.6	0.4	6.1	5.8	2016
ES LOVELY	34.4	17.5	99	11.40	200	34.1	5.98	56.5	7.5	8.3	0.1	1.5	6.0	3.5	2016
ROGOSO	29.1	17.5	98	11.02	192	29.1	5.07	55.8	6.5	8.3	0.0	0.0	6.8	6.3	2016
ES REGAIN	34.2	17.3	98	11.45	199	33.1	5.75	57.1	7.3	8.3	0.2	0.4	6.3	3.7	2007
ARCADE	34.2	17.2	97	11.62	200	34.0	5.86	58.5	7.0	8.3	0.3	0.5	6.7	4.8	2012
ES REMINGTON	33.3	17.1	96	11.17	191	31.9	5.46	55.7	7.4	8.3	0.3	0.3	6.5	4.1	2014
RGT EARLEXX	32.9	16.9	95	11.57	195	32.3	5.45	58.4	6.8	8.2	0.8	0.7	6.3	3.3	2017
ES ARDENT	35.3	16.1	91	11.35	182	34.1	5.47	56.1	6.5	8.3	0.4	0.5	6.0	5.7	2011
Trial statistics															
Mean of 4 & 5 year varieties	32.7	17.7	100	11.48	204	32.3	5.72	57.8	7.2	8.2	0.6	0.4	6.8	4.9	-
LSD (V.Cont)	0.82	0.46	2.6	0.123	-	1.34	-	0.64	0.29	-	-	-	0.45	-	-
SE (variety mean)	0.412	0.234	1.32	0.0618	-	0.678	-	0.325	0.145	-	-	-	0.226	-	-

On the 1-9 scales, high figures indicate that a variety shows the character to a high degree (e.g. high resistance)

Notes:
 ME is calculated as D-value x 0.16
 # = values are re-transformed from the mean of the log data
 Limited data for brackling % with some varieties only having 1 trials worth of data
 ** = Eyespot rating derived from inoculated nurseries rather than field scores



With thanks to the Maize Growers Association for part-funding the eyespot assessments



2019 FORAGE MAIZE DESCRIPTIVE LIST

First choice varieties for less favourable sites (varieties ranked in dry matter yield order)

VARIETY	Dry matter content at harvest (%)	Dry matter yield (t/ha)	Dry matter yield (% of 4 and 5 year varieties)	ME of fresh plant at harvest (MJ/kg dry matter)	ME yield of fresh plant at harvest (1000's MJ/ha)	Starch content of whole plant at harvest (%)	Starch yield of whole plant at harvest (t/ha)	Cell wall digestibility (%)	Early vigour (1-9)	Standing power at harvest (root lodging) (1-9)	Lodging (%) #	Brackling (%) #	Leaf senescence (1-9)	Eyespot rating (1-9) **	Year first listed
EMMERSON	37.6	16.3	95	11.44	187	35.3	5.76	56.5	7.0	8.3	0.3	1.1	4.9	2.6	2015
CITO KWS	37.3	16.1	94	11.61	187	36.0	5.81	57.7	7.3	8.3	0.2	1.5	4.4	5.4	2018
AUGUSTUS KWS	36.2	16.3	95	11.49	188	35.2	5.74	57.0	7.2	8.1	1.7	2.4	4.7	6.0	2015
RUBIERA KWS	36.0	16.4	95	11.62	190	36.2	5.92	57.7	7.4	8.1	2.3	2.0	4.7	5.4	2015
SERGIO KWS	35.3	16.5	96	11.56	191	36.1	5.95	57.4	7.5	8.0	3.0	0.6	5.2	5.7	2014
REASON	35.3	17.6	102	11.44	201	32.6	5.75	57.2	7.0	8.2	1.1	0.8	5.9	3.4	2016
AVITUS KWS	35.2	18.4	107	11.31	208	34.3	6.32	55.7	7.7	8.2	1.2	3.3	5.2	5.7	2018
ACTIVATE	35.2	16.5	96	11.65	192	35.3	5.82	58.3	6.9	8.3	0.3	0.0	6.5	4.6	2012
PEREZ KWS	34.4	17.5	102	11.24	197	33.3	5.85	55.8	7.7	8.1	1.7	0.7	5.7	4.8	2015
GLORY	34.3	17.4	101	11.41	198	34.1	5.93	56.8	7.3	8.3	0.3	0.2	6.7	6.3	2013
YUKON	33.3	17.4	101	11.22	195	34.0	5.93	55.2	7.1	8.3	0.3	0.2	6.5	5.0	2012
AMBITION	33.2	17.8	103	11.40	202	32.9	5.85	57.1	7.4	8.3	0.2	0.6	6.8	6.3	2012
MADONIAS	33.1	17.5	102	11.34	198	33.5	5.86	56.4	7.2	8.2	0.4	2.9	6.0	5.5	2018
SUNLITE	33.0	16.9	98	11.57	195	33.8	5.70	58.6	7.1	8.3	0.2	0.0	6.5	3.0	2014
SEVERUS	33.0	17.4	101	11.27	196	33.2	5.76	56.1	7.3	8.2	0.5	0.3	6.2	5.3	2013
PINNACLE	32.9	17.4	101	11.50	201	34.3	5.98	57.3	7.3	8.2	1.1	2.3	7.2	6.6	2018
AUTENS KWS	32.9	18.0	105	11.22	202	31.6	5.69	56.0	7.4	8.0	2.4	0.4	6.0	5.0	2017
AURELIUS KWS	32.4	18.1	105	11.24	203	31.5	5.69	56.2	7.7	8.1	1.5	1.4	6.3	4.6	2016
FIELDSTAR	32.1	17.9	104	11.35	203	32.0	5.72	57.2	7.5	8.3	0.2	0.2	7.3	6.5	2013
RODRIGUEZ KWS	31.8	17.6	102	11.42	201	33.8	5.94	57.4	6.8	8.3	0.1	1.0	6.9	3.8	2015
BONNIE	31.0	17.8	103	11.48	205	30.7	5.47	58.7	7.3	8.2	0.7	0.1	7.6	5.8	2017
GATSBY	30.9	17.8	103	11.31	202	30.4	5.41	57.2	7.3	8.3	0.1	0.1	7.4	5.4	2017
EMBLEM	30.9	17.5	101	11.34	198	32.0	5.59	57.6	7.4	8.3	0.4	0.1	6.8	5.7	2014
ABSALON	30.9	17.9	104	11.45	205	29.1	5.19	59.1	7.0	8.3	0.4	0.0	7.3	5.9	2017
ASGAARD	30.6	17.6	102	11.36	200	29.8	5.24	58.2	7.6	8.2	0.4	0.0	7.3	3.8	2014
BELAMI CS	30.0	17.5	102	11.47	201	29.4	5.14	59.2	7.3	7.9	3.8	0.0	6.8	4.5	2017
Trial statistics															
Mean of 4 & 5 year varieties	33.1	17.2	100	11.37	196	32.5	5.60	57.2	7.2	8.2	0.7	0.5	6.4	-	-
LSD (V.Cont)	1.12	0.54	3.2	0.157	-	1.68	-	0.82	0.37	-	-	-	0.56	-	-
SE (variety mean)	0.564	0.272	1.58	0.0789	-	0.840	-	0.412	0.186	-	-	-	0.283	-	-

On the 1-9 scales, high figures indicate that a variety shows the character to a high degree (e.g. high resistance)

Notes:

- = no data available

ME is calculated as D-value x 0.16

= values are re-transformed from the mean of the log data

Limited data for brackling % with some varieties only having 1 trials worth of data so 2012 trials still included

** = Eyespot rating derived from inoculated nurseries rather than field scores



With thanks to the Maize Growers Association for part-funding the eyespot assessments

2019 FORAGE MAIZE DESCRIPTIVE LIST

Second choice varieties for less favourable sites (varieties ranked in dry matter yield order)

VARIETY	Dry matter content at harvest (%)	Dry matter yield (t/ha)	Dry matter yield (% of 4 and 5 year varieties)	ME of fresh plant at harvest (MJ/kg dry matter)	ME yield of fresh plant at harvest (1000's MJ/ha)	Starch content of whole plant at harvest (%)	Starch yield of whole plant at harvest (t/ha)	Cell wall digestibility (%)	Early vigour (1-9)	Standing power at harvest (root lodging) (1-9)	Lodging (%) #	Brackling (%) #	Leaf senescence (1-9)	Eyespot rating (1-9) **	Year first listed
ES ARDENT	35.5	15.8	92	11.22	178	33.5	5.30	55.6	6.4	8.3	0.3	1.4	5.4	5.7	2011
ES REGAIN	33.7	17.1	99	11.32	193	32.4	5.54	56.7	7.5	8.3	0.3	0.9	6.2	3.7	2007
ARCADE	33.5	16.9	98	11.48	194	33.0	5.59	58.0	6.8	8.2	0.4	1.4	6.6	4.8	2012
ES LOVELY	33.5	17.0	99	11.19	191	32.8	5.58	55.8	7.4	8.3	0.1	1.0	6.1	3.5	2016
RGT OXXGOOD	32.8	17.4	101	11.29	197	31.9	5.56	56.9	7.1	8.2	0.8	0.8	6.2	5.8	2016
ES REMINGTON	32.6	17.0	99	11.01	187	30.6	5.20	55.3	7.3	8.2	0.5	0.3	6.6	4.1	2014
RGT EARLEXX	32.1	16.4	95	11.38	187	31.0	5.08	57.7	6.8	8.2	0.7	1.4	6.3	3.3	2017
MONTY	31.3	17.9	104	11.31	203	29.0	5.20	57.8	7.0	8.3	0.3	0.5	7.6	4.5	2014
SY NORDICSTAR	31.0	17.5	101	11.27	197	30.5	5.32	57.8	7.4	8.1	1.4	0.3	7.5	7.5	2016
MILKSTAR	29.7	18.4	107	10.86	200	22.9	4.21	56.3	7.3	7.9	3.8	0.3	7.4	5.4	2018
Trial statistics															
Mean of 4 & 5 year varieties	33.1	17.2	100	11.37	196	32.5	5.60	57.2	7.2	8.2	0.7	0.5	6.4	-	-
LSD (V.Cont)	1.12	0.54	3.2	0.157	-	1.68	-	0.82	0.37	-	-	-	0.56	-	-
SE (variety mean)	0.564	0.272	1.58	0.0789	-	0.840	-	0.412	0.186	-	-	-	0.283	-	-

On the 1-9 scales, high figures indicate that a variety shows the character to a high degree (e.g. high resistance)

Notes:

- = no data available

ME is calculated as D-value x 0.16

= values are re-transformed from the mean of the log data

Limited data for brackling % with some varieties only having 1 trials worth of data so 2012 trials still included

** = Eyespot rating derived from inoculated nurseries rather than field scores



With thanks to the Maize Growers Association for part-funding the eyespot assessments

FORAGE MAIZE FOR ANAEROBIC DIGESTION

Varieties for favourable sites (2019)

VARIETY	Dry matter content at harvest (%)	Dry matter yield (t/ha)	Dry matter yield (% of 4 and 5 year varieties)	ME of fresh plant at harvest (MJ/kg dry matter)	ME yield of fresh plant at harvest (1000's MJ/ha)	Early vigour (1-9)	Standing power at harvest (root lodging) (1-9)	Eyespot rating (1-9)
LG31218	30.5	18.7	105	11.60	217	7.6	8.0	5.4
LG31211	30.3	18.5	105	11.60	215	7.6	8.0	4.6
BONNIE	32.3	18.2	103	11.66	213	7.2	8.3	5.8
GATSBY	31.8	18.4	104	11.51	212	7.2	8.3	5.4
LG30209	30.9	18.3	103	11.54	212	7.5	8.1	3.3
ATRIUM	30.4	18.0	102	11.70	210	7.4	8.2	2.8
CATHY	30.4	18.5	105	11.34	210	7.2	8.3	4.8
ABSALON	31.5	18.2	103	11.53	210	7.1	8.3	5.9
MONTY	32.2	18.1	102	11.47	207	6.9	8.3	4.5
ASGAARD	32.0	18.0	101	11.51	207	7.7	8.3	3.8
FIELDSTAR	33.0	17.9	101	11.54	206	7.3	8.3	6.5
TIBERIO	29.6	18.4	104	11.20	206	7.4	8.1	6.3
SY NORDICSTAR	32.3	17.9	101	11.50	205	7.4	8.2	7.5
CODITANK	29.7	18.0	101	11.30	203	6.5	8.3	5.0
ES MARCO	29.3	18.1	102	11.18	202	6.4	8.2	6.9
ES LOVELY	34.4	17.5	99	11.40	200	7.5	8.3	3.5
ES REGAIN	34.2	17.3	98	11.45	199	7.3	8.3	3.7
MAS 13M	30.1	18.3	103	10.86	199	6.8	8.2	5.2
ROGOSO	29.1	17.5	98	11.02	192	6.5	8.3	6.3
ES REMINGTON	33.3	17.1	96	11.17	191	7.4	8.3	4.1
EMMERSON	37.6	16.4	93	11.62	191	7.0	8.3	2.6
ES ARDENT	35.3	16.1	91	11.35	182	6.5	8.3	5.7
Trial statistics								
Mean of 4 & 5 year varieties	32.7	17.7	100	11.48	204	7.2	8.2	4.9
LSD (V.Cont)	0.82	0.46	2.6	0.123	-	0.29	-	-
SE (variety mean)	0.412	0.234	1.32	0.0618	-	0.145	-	-

On the 1-9 scales, high figures indicate that a variety shows the character to a high degree (e.g. high resistance)

Notes:

ME is calculated as D-value x 0.16

** = Eyespot rating derived from inoculated nurseries rather than field scores

The varieties tabulated are forage maize varieties that the breeders have nominated as having potential suitability for AD use. The List may not represent the full range of forage maize or biogas varieties that are available for the market.



With thanks to the Maize Growers Association for part-funding the eyespot assessments

FORAGE MAIZE FOR ANAEROBIC DIGESTION

Varieties for less favourable sites (2019)

VARIETY	Dry matter content at harvest (%)	Dry matter yield (t/ha)	Dry matter yield (% of 4 and 5 year varieties)	ME of fresh plant at harvest (MJ/kg dry matter)	ME yield of fresh plant at harvest (1000's MJ/ha)	Early vigour (1-9)	Standing power at harvest (root lodging) (1-9)	Eyespot rating (1-9)
EMMERSON	37.6	16.3	95	11.44	187	7.0	8.3	2.6
ES ARDENT	35.5	15.8	92	11.22	178	6.4	8.3	5.7
ES REGAIN	33.7	17.1	99	11.32	193	7.5	8.3	3.7
ES LOVELY	33.5	17.0	99	11.19	191	7.4	8.3	3.5
ES REMINGTON	32.6	17.0	99	11.01	187	7.3	8.2	4.1
MONTY	31.3	17.9	104	11.31	203	7.0	8.3	4.5
SY NORDICSTAR	31.0	17.5	101	11.27	197	7.4	8.1	7.5
FIELDSTAR	32.1	17.9	104	11.35	203	7.5	8.3	6.5
BONNIE	31.0	17.8	103	11.48	205	7.3	8.2	5.8
GATSBY	30.9	17.8	103	11.31	202	7.3	8.3	5.4
ABSALON	30.9	17.9	104	11.45	205	7.0	8.3	5.9
ASGAARD	30.6	17.6	102	11.36	200	7.6	8.2	3.8
Trial statistics								
Mean of 4 & 5 year varieties	33.1	17.2	100	11.37	196	7.2	8.2	4.9
LSD (V.Cont)	1.12	0.54	3.2	0.157	-	0.37	-	-
SE (variety mean)	0.564	0.272	1.58	0.0789	-	0.186	-	-

On the 1-9 scales, high figures indicate that a variety shows the character to a high degree (e.g. high resistance)

Notes:

ME is calculated as D-value x 0.16

** = Eyespot rating derived from inoculated nurseries rather than field scores

The varieties tabulated are forage maize varieties that the breeders have nominated as having potential suitability for AD use. The List may not represent the full range of forage maize or biogas varieties that are available for the market.



With thanks to the Maize Growers Association for part-funding the eyespot assessments